

*Lill* *fm*

National Aeronautics and Space Administration  
Goddard Space Flight Center  
Contract No. NAS-5-3760

ST - GM - SP - 10401

VARIATION OF SOLAR WIND INTENSITY WITH THE PHASE OF  
THE SOLAR ACTIVITY CYCLE ACCORDING TO DATA OF  
STEADY GEOMAGNETIC FIELD OSCILLATIONS

by

O. V. Bol'shakova  
[USSR]

FACILITY FORM 502

**N66-87176**

(ACCESSION NUMBER)

(PAGES)

**CR 78207**  
(NASA CR OR TMX OR AD NUMBER)

(THRU)

(CODE)

(CATEGORY)

15 OCTOBER 1965

VARIATION OF SOLAR WIND INTENSITY WITH THE PHASE OF  
THE SOLAR ACTIVITY CYCLE ACCORDING TO DATA OF  
STEADY GEOMAGNETIC FIELD OSCILLATIONS\*

Astronomicheskiy Zhurnal  
Tom 42, vyp. 4, 859 - 861,  
Izdatel'stvo "NAUKA", 1965.

by O. V. Bol'shakova

SUMMARY

It is shown in this work that the mean amplitude of short-period oscillations of the Earth's magnetic field of the type pc<sup>4</sup>, corresponding to the solar wind from unperturbed regions of the Sun ( $K_p = 0$ ), remained constant during the period from 1957 through 1964. Hence the conclusion is derived that, according to [1], the intensity of the solar wind originating from the unperturbed regions of the Sun does not vary with the phase of solar activity cycle.

\*  
\*   \*  
\*

It was shown in the work by M. S. Bobrov et al [1] that the properties of the solar wind are investigated on the basis of study of permanent variations in polar caps ( $P_o$ - disturbances) during days with a very low planetary magnetic activity. One of the results of the work [1] is the conclusion that the solar wind is continually generated by the quiet Sun regions, and that the emissive capability of those regions does not depend on the level of solar activity.

It appears to be quite important to investigate this question on the basis of quite independent data on short-period oscillations of the Earth's electromagnetic field. It is well known from short-period oscillations' morphology that there exists a dependence of the period of steady

---

\* IZMENENIYE INTENSIVNOSTI SOLNECHNOGO VETRA S FAZOY TSIKLA SOLNECHNOY ATIVNOSTI PO DANNYM USTOYCHIVYKH KOLEBANIY GEOMAGNITNOGO POLYA.

oscillations of the geomagnetic field on the level of magnetic activity, so that to each level of activity corresponds a specific group of oscillations. In particular, to the zero level of activity corresponds the type- $pc4$  group of oscillations with periods from 50 to 150 sec. [2]. The  $pc4$  - oscillations are observed at low, practically zero magnetic activity, and consequently they cannot be identified with corpuscular fluxes from active regions and flares. This, and also the presence in the given group of oscillations of entirely specific daily and seasonal course allows to make the assumption that  $pc4$  are consequence of the action upon the magnetosphere of the Earth of only the solar wind from unperturbed regions of the Sun.

It should be noted that  $pc4$ -oscillations are observed practically on every day corresponding to a low planetary activity, and their amplitude varies from zero to a certain maximum value (of the order of hundredths of fractions of gamma for the vertical component).

Therefore, the variation of solar wind intensity with the phase of solar activity cycle may be detected by merely studying the seasonal course of the amplitude of  $pc4$ , averaged precisely in the same way as was done in the work [1] relative to amplitudes of the H- component. A significant time interval, encompassing a very high activity (1957 - 1959) was investigated alongside with the activity drop (1960 - 1961) and the low solar activity of 1961 - 1964 according to the variations of the vertical and horizontal components of the geomagnetic field at st. Borok. During that period,  $pc4$  - oscillations were observed for 267 days with fairly low magnetic activity, that is, practically always when recurrent or sporadic disturbances were absent, and when only the corpuscular radiation from the unperturbed regions of the Sun acted upon the Earth's magnetosphere.

We plotted in Fig. 1 [next page] in ordinates the amplitudes of  $pc4$  in gammas, averaged at the beginning for all hours of the given day whenever  $pc4$  were observed, and then for the season by all the days for which  $\sum K_p \leq 10$ . As may be seen from the graph, the amplitude of  $pc4$  has a clearly marked seasonal course with a summer maximum and winter minimum, and the value of the wintertime as well as summertime extremes of amplitude does not practically change with the phase of the solar activity cycle.

The exact correspondence of the extremes on curve on Fig. 1 for summer and winter points to the absence of the influence of equinoctial maxima of

magnetic activity due to recurrent perturbations. This influence is manifest in "creeping" of amplitude maxima of the seasonal course for the oscillations of the pc3-type group corresponding to recurrent disturbances; at the same time the mean amplitude of pc3 drops in the cycle of solar activity from the maximum to the minimum epoch by about a factor of 2 [2]. The fact of constance of extremum position in the seasonal course of the pc4 amplitude is further evidence in favor of the fact that pc4 correspond precisely to the solar wind originating in the unperturbed regions of the Sun.




Fig. 1. - Seasonal course of the amplitude of pc4 in a cycle of solar activity at the st. Borok in 1957-1964. The ordinates represent the amplitude of the vertical component of pc4 in gammas.

Therefore, in magnetoquiet days ( $K_p \approx 0$ ) the solar wind does not modify its intensity even according to data of short-period oscillations of the Earth's, as a function of the phase of the solar activity cycle; it constitutes the corpuscular radiation from unperturbed regions of the Sun.

\*\*\* THE END \*\*\*

Geophysical Station Borok  
of the  
Institute of Earth's Physics  
of the  
USSR Academy of Sciences

Received on 29 March 1965.

Contract No. NAS-5-3760  
Consultants & Designers, Inc.  
Arlington, Virginia

Translated by ANDRE L. BRICHANT  
on 14 October 1965

REFERENCES

- [1].- M. S. BOBROV, N. F. KOROLEVA, R. M. NOVIKOVA.- Geomagn. i Aeron. 4, 333,  
 [2].- V. A. TROITSKAYA.- Research in Geophys. 1, 485, 1964 1964.

DISTRIBUTIONGODDARD SPACE F.C.

100 CLARK, TOWNSEND  
 110 STROUD  
 400 BOURDEAU  
 600 PIEPER  
 610 MEREDITH  
       SEDDON  
 611 McDONALD  
       L. DAVIS  
       CLINE  
       ABRAHAM  
       BOLDT  
 612 HEPPNER  
       NESS  
       SUGIURA  
 613 KUPPERIAN [3]  
 614  
       WHITE  
       HALLAM  
 615 BAUER  
       WHIPPLE  
       GOLDBERG  
       SERBU  
       STONE  
 640 HESS [3]  
       MEAD  
       NAKADA  
       SPEISER  
       STERN  
 630 GI for SS [5]  
 620 SPENCER  
       NEWTON  
 252 LIBRARY  
 256 FREAS

NASA HQS

SS NEWELL, NAUGLE  
 SG MITCHELL  
       SCHARDT  
       SMITH  
       DUBIN  
 SL LIDDEL  
       FELLOWS  
       HIPSHER  
       HOROWITZ  
 SM FOSTER  
       ALLENBY  
       GILL  
       BADGLEY  
 RR KURZWEG  
 RRP GESSOW  
 RV-1 PEARSON  
 RTR NEILL  
 ATSS SCHWIND [2]  
       ROBBINS  
 WX SWEET

OTHER CENTERSARC

SONETT [5]  
 LIBRARY [3]

LaRC

160 ADAMSON  
       HESS  
 185 WEATHERWAX 2  
 213 KATZOFF

JPL

SNYDER  
 NEUGEBAUER

UCLA

COLEMAN

UC BERKELEY

WILCOX